

Temperature dependence of the response function of hot nuclear matter

Abdellatif Abada¹ and Dominique Vautherin²

¹ *Theoretical Physics Group, Department of Physics and Astronomy,
University of Manchester, Manchester M13 9PL, UK*

² *Division de Physique Théorique *, Institut de Physique Nucléaire
Orsay Cedex, F-91406, France*

ABSTRACT

The description of collective motion in nuclei at finite temperature using the framework of the random phase approximation is discussed. We focus on the special case of the isovector response function of hot nuclear matter using various effective Skyrme interactions.

MC/TH 95/20

1. Introduction

The purpose of the present notes is to investigate the response function of hot nuclear matter. We use the framework of time dependent mean field theory with the standard Skyrme type effective interactions which have provided in the past good descriptions of average nuclear properties. For such interactions analytic expressions for the response function are derived. From these analytic formulae we show that one can gain an understanding of the temperature dependance of the giant dipole resonance in finite nuclei. Although effects which are beyond the framework of linear response theory are important to account for the full width of the giant dipole state (e.g. two particle two hole configurations) many features of these resonances are known to be readily understood at this level already ^{1,2,3,4,5}. In the following we show that zero sound type modes become less collective as temperature rises and sometimes disappear at temperatures of a few MeV. We also show that for effective interactions with an effective mass close to unity collective effects are small already at zero temperature and depend weakly on temperature. We will thus argue that the temperature dependance of the giant dipole resonance provides valuable information on the effective interaction.

2. Linear response theory in hot nuclear matter

When an external field of the form

$$V_{\text{ext}} = \epsilon \tau_3 e^{i(\mathbf{q} \cdot \mathbf{r} - (\omega + i\eta)t)}, \quad (1)$$

*Unité de Recherche des Universités Paris XI et Paris VI associée au CNRS

is applied to nuclear matter, it induces a density change between neutron and proton densities which is of the form

$$\delta\rho(\mathbf{r}, t) = \rho_n(\mathbf{r}, t) - \rho(\mathbf{r}, t) = \eta e^{i(\mathbf{q}\cdot\mathbf{r} - (\omega + i\eta)t)}. \quad (2)$$

The ratio of the amplitude α of the density oscillations to the external field strength is the response function (also called dynamic polarizability)

$$\Pi(\omega, q) = \frac{\alpha}{\epsilon} \quad (3)$$

In the case of Skyrme type forces it is possible to find analytic expressions for the response function of nuclear matter at finite temperature^{6,7}.

Calculations are particularly simple in the case of a simplified Skyrme interaction

$$v = t_0(1 + x_0 P_\sigma)\delta(\mathbf{r}_1 - \mathbf{r}_2) + t_3\delta(\mathbf{r}_1 - \mathbf{r}_2)\delta(\mathbf{r}_2 - \mathbf{r}_3). \quad (4)$$

In this particular case one has the simple formula

$$\Pi(\omega, \mathbf{q}) = \frac{\Pi_0(\omega, \mathbf{q})}{1 - V_0\Pi_0(\omega, \mathbf{q})}, \quad (5)$$

where Π_0 is the unperturbed polarization propagator⁸, also called Lindhard function

$$\Pi_0(\omega, \mathbf{q}) = \frac{2}{(2\pi)^3} \int d\mathbf{k} \frac{f(\mathbf{k} + \mathbf{q}) - f(\mathbf{k})}{\omega + i\eta - \epsilon(\mathbf{k}) + \epsilon(\mathbf{k} + \mathbf{q})}. \quad (6)$$

and where

$$V_0 = -t_0(x_0 + 1/2)/2 - t_3\rho_0/8. \quad (7)$$

For a given momentum $\mathbf{q}$ we thus see that there is a resonant response when the frequency ω of the external field corresponds to a zero in the denominator i.e. when

$$1 = V_0\Pi_0(\omega, \mathbf{q}). \quad (8)$$

The real part of ω determines the energy of the collective mode and its imaginary part the lifetime⁸.

The previous formulae correspond to the special case of a simplified Skyrme force. They can however be generalized to the case of a full Skyrme force. Such an interaction contains a density independent part v_{12} and a density dependent part v_{123} . The density independent part is parametrized in momentum space as

$$\langle \mathbf{k} | v_{12} | \mathbf{k}' \rangle = t_0(1 + x_0 P_\sigma) + \frac{t_1}{2}(1 + x_1 P_\sigma)(k^2 + k'^2) + t_2(1 + x_2 P_\sigma)\mathbf{k}\cdot\mathbf{k}'. \quad (9)$$

The density dependent part in ordinary space is

$$v_{123} = \frac{t_3}{6}(1 + x_3 P_\sigma)\delta(\mathbf{r}_1 - \mathbf{r}_2)\rho^\sigma\left(\frac{\mathbf{r}_1 + \mathbf{r}_2}{2}\right). \quad (10)$$

For such an interaction the response function is given by

$$\Pi(\omega, \mathbf{q}) = \frac{\Pi_0(\omega, \mathbf{q})}{1 - \bar{V}_0\Pi_0(\omega, \mathbf{q}) - 2V_1\Pi_2(\omega, \mathbf{q}) - V_1^2\Pi_2^2(\omega, \mathbf{q}) - V_1^2\Pi_4\Pi_0}. \quad (11)$$

In this equation Π_2 and Π_4 are generalized Lindhard functions defined by

$$\Pi_{2N}(\omega, \mathbf{q}) = \frac{4}{(2\pi)^3} \int d^3k \frac{f(\mathbf{k} + \mathbf{q}) - f(\mathbf{k})}{\omega + i\eta - \epsilon(\mathbf{k}) + \epsilon(\mathbf{k} + \mathbf{q})} (\mathbf{k} \cdot (\mathbf{k} + \mathbf{q}))^N, \quad (12)$$

while V_0 and V_2 are linear combinations of the parameters of the Skyrme interaction. Explicitly one has

$$\begin{aligned} V_0 = & -\frac{t_0}{2} \left(x_0 + \frac{1}{2} \right) - \frac{t_3}{12} \left(x_3 + \frac{1}{2} \right) \rho_0^\alpha \\ & - \frac{q^2}{16} (3t_1(1 + 2x_1) + t_2(1 + 2x_2)) \\ & - \left(\frac{m^*\omega}{q} \right)^2 \frac{2V_1}{1 - 2V_1m^*\rho_0}, \end{aligned} \quad (13)$$

where m^* is the effective mass and ρ_0 the saturation density of nuclear matter, while V_1 is:

$$V_1 = \frac{1}{16} (t_2(1 + 2x_2) - t_1(1 + 2x_1)) \quad (14)$$

In reference ¹⁰ a discussion of the properties of the response function in the case of a simplified Skyrme interaction was provided. It was found that for a momentum transfer $q=0.23 \text{ fm}^{-1}$ (a value which corresponds to the giant dipole resonance in lead- 208 in the Steinwedel- Jenssen model) the response function exhibits a sharp zero sound type peak at zero temperature. In contrast when temperature rises the smearing of the Lindhard function was found to make the peak much weaker and in fact nearly disappear for temperatures of a few MeV. Whether this conclusion holds for the more realistic Skyrme forces is a question which was investigated in reference ⁶. Some results of this study are shown in Figure 1.

Fig.1a (left side) gives the strength

$$S(\omega) = -\frac{1}{\pi} \Im m \Pi(\omega, q). \quad (15)$$

as a function of the frequency ω for a momentum transfer $q= 0.23 \text{ fm}^{-1}$ in the case of the Skyrme force SIII while Fig.1b (right side) corresponds to interaction SV. It can be seen that for interaction SIII collective effects disappear as soon as temperature reaches a few MeV while for SV only a weak temperature dependance is observed. In reference ⁶ the difference between these two cases was attributed to a difference between the effective mass values for the two interactions. This difference shows that the temperature dependance of the response function provides information on the effective particle hole interaction. In this context it is worthwhile noting




Fig. 1. Distribution of strength as a function of the frequency ω for a momentum transfer $q=0.23 \text{ fm}^{-1}$ in the case of interactions SIII and SV

that recent experimental studies of the giant dipole resonance found a saturation of photon multiplicities at high excitation energy¹¹. This may be a signature of the loss of collectivity found at high temperature in the present calculation.

Acknowledgements One of us (D.V.) wishes to thank Prof. Apolodor Raduta for his invitation to present this lecture at the 1995 Predeal School on Nuclear Structure.

1. M. E. Faber, J. L. Egido and P. Ring, Phys. Lett. **127B** (1983) 5
2. M. Sommerman, Ann. of Phys. **151** (1983) 163
3. H. Sagawa and G. F. Bertsch, Phys. Lett. **333B** (1994) 289
4. D. Vautherin and N. Vinh Mau, Nucl. Phys. **A422** (1984) 140
5. N. Vinh Mau, Nucl. Phys. **A548** (1992) 381
6. F. Braghin, D. Vautherin and A. Abada, Phys. Rev. C (1995), in press
7. E.S.Hernandez, J. Navarro, A. Polls and J. Ventura, Finite Temperature RPA in Symmetric Nuclear Matter with Skyrme Interactions, University of Valencia preprint, July 27th, 1995
8. A.L. Fetter and J.D. Walecka, Quantum Theory of Many Particle Systems (Mc Graw-Hill, New York, 1971)
9. C. Garcia- Recio, J. Navarro, Nguyen van Giai and L.L Salcedo Ann. Phys. **214** (1992) 293
10. F. Braghin and D. Vautherin, Phys. Lett. **333B** (1994) 289
11. J. H. Le Faou *et al.*, Phys. Rev. Lett. **72** (1994) 3321